



## Decision Analysis

Publication details, including instructions for authors and subscription information:  
<http://pubsonline.informs.org>

### Quantile-Parameterized Distributions for Expert Knowledge Elicitation

Dmytro Perepolkin, Erik Lindström, Ullrika Sahlin

To cite this article:

Dmytro Perepolkin, Erik Lindström, Ullrika Sahlin (2025) Quantile-Parameterized Distributions for Expert Knowledge Elicitation. *Decision Analysis* 22(3):169-188. <https://doi.org/10.1287/deca.2024.0219>

This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License. You are free to download this work and share with others for any purpose, even commercially if you distribute your contributions under the same license as the original, and you must attribute this work as “*Decision Analysis*.” Copyright © 2025 The Author(s). <https://doi.org/10.1287/deca.2024.0219>, used under a Creative Commons Attribution License: <https://creativecommons.org/licenses/by-sa/4.0/>.”

Copyright © 2025 The Author(s)

Please scroll down for article—it is on subsequent pages



With 12,500 members from nearly 90 countries, INFORMS is the largest international association of operations research (O.R.) and analytics professionals and students. INFORMS provides unique networking and learning opportunities for individual professionals, and organizations of all types and sizes, to better understand and use O.R. and analytics tools and methods to transform strategic visions and achieve better outcomes.

For more information on INFORMS, its publications, membership, or meetings visit <http://www.informs.org>

# Quantile-Parameterized Distributions for Expert Knowledge Elicitation

Dmytro Perepolkin,<sup>a,\*</sup> Erik Lindström,<sup>a</sup> Ullrika Sahlin<sup>b</sup>

<sup>a</sup> Centre for Mathematical Sciences, Lund University, 223 62 Lund, Sweden; <sup>b</sup> Centre for Environmental and Climate Science, Lund University, 223 62 Lund, Sweden

\*Corresponding author

Contact: [dmytro.perepolkin@matstat.lu.se](mailto:dmytro.perepolkin@matstat.lu.se),  <https://orcid.org/0000-0001-8558-6183> (DP); [erik.lindstrom@matstat.lu.se](mailto:erik.lindstrom@matstat.lu.se),  <https://orcid.org/0000-0002-6468-2624> (EL); [ullrika.sahlin@cec.lu.se](mailto:ullrika.sahlin@cec.lu.se),  <https://orcid.org/0000-0002-2932-6253> (US)

Received: July 3, 2024

Revised: September 24, 2024


Accepted: December 25, 2024

Published Online in Articles in Advance:  
March 31, 2025

<https://doi.org/10.1287/deca.2024.0219>

Copyright: © 2025 The Author(s)

**Abstract.** This paper provides a comprehensive overview of quantile-parameterized distributions (QPDs) as a tool for capturing expert predictions and parametric judgments. We survey a range of methods for constructing distributions that are parameterized by a set of quantile-probability pairs and describe an approach to generalizing them to enhance their tail flexibility. Furthermore, we explore the extension of QPDs to the multivariate setting, surveying the approaches to construct bivariate distributions, which can be adopted to obtain distributions with quantile-parameterized margins. Through this review and synthesis of the previously proposed methods, we aim to enhance the understanding and utilization of QPDs in various domains.

 **Open Access Statement:** This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License. You are free to download this work and share with others for any purpose, even commercially if you distribute your contributions under the same license as the original, and you must attribute this work as “*Decision Analysis*. Copyright © 2025 The Author(s). <https://doi.org/10.1287/deca.2024.0219>, used under a Creative Commons Attribution License: <https://creativecommons.org/licenses/by-sa/4.0/>.”

**Funding:** U. Sahlin was funded by the Crafoord Foundation [ref 20200626].

**Supplemental Material:** The online appendix is available at <https://doi.org/10.1287/deca.2024.0219>.

**Keywords:** quantile functions • quantile-parameterized distributions • expert knowledge elicitation • Bayesian analysis

## 1. Introduction

Judgment plays a crucial role in transforming raw data into meaningful insights. To be useful, judgment must be translated into mathematical models and assumptions. These models are designed to capture the expert’s understanding of the world, including the causal links between relevant entities. The models serve as a representation of this understanding while also addressing knowledge limitations, treated as uncertainties. Elicitation involves translating qualitative understanding into quantitative models offering valuable insights.

Most of the expert elicitation protocols described in the literature (Spetzler and Staël Von Holstein 1975, O’Hagan et al. 2006, Morgan 2014, Gosling 2018, Hemming et al. 2018, Welsh and Begg 2018, Hanea et al. 2021) encode

expert judgments about the parameter or quantity of interest as an ordered set of quantiles with corresponding probabilities. This typically includes measures such as the median and the upper and lower quartiles. Assessors are then encouraged to select a probability distribution that reasonably fits the elicited quantile-probability pairs and validate the choice with the expert (Gosling 2018). A distribution is selected from a predefined set of “simple and convenient” distributions (O’Hagan et al. 2006) with boundedness that accounts for the nature of the elicited quantity.

Several specialized distributions have been developed to facilitate smooth interpolation of probabilistic assessments. These distributions, parameterized by quantile-probability pairs, ensure that the elicited quantile-probability pairs (QPPs) are preserved exactly

(Keelin and Powley 2011, Powley 2013, Keelin 2016, Hadlock 2017, Wilson et al. 2023). Quantile-parameterized distributions are particularly valuable thanks to the interpretability of their parameters. By leveraging the elicited quantiles, these distributions enable precise capturing of expert knowledge while maintaining a high level of flexibility in modeling.

We believe that the primary utility of QPDs lies in their ability to simplify the specification of probability distributions for model parameters, known as *prior elicitation* (Mikkola et al. 2023). However, these same distributions can also be employed to describe an expert's predictions for the next observation, referred to as *predictive elicitation* (Kadane 1980, Winkler 1980, Akbarov 2009, Hartmann et al. 2020), or to capture both uncertainty and variability through a two-dimensional probability distribution in *hybrid elicitation* (Perepolkin et al. 2024).

With this work, we aim to introduce quantile-parameterized distributions (QPDs) to a wide readership. The literature review and the findings are presented through the perspective of quantile functions, building upon the theoretical foundations established by Parzen (1979) and Gilchrist (2000). The derivatives and inverses for each of the quantile functions discussed in this paper are provided in the Supplementary Materials, serving as a valuable reference for future research. Through our comprehensive review and identification of research gaps, we aim to contribute to the development of flexible and extensible distributions that can effectively capture expert knowledge.

### 1.1. Paper Structure

In Section 2, we revisit the approaches to quantile parameterization of probability distributions and explore how QPDs can effectively describe expert beliefs regarding model parameters or predictions. In Section 3, we conduct a comprehensive review and comparison of various continuous univariate QPDs found in the literature. Specifically, we focus on the Myerson distribution and its generalization accommodating different tail thicknesses. We compare the robust moments of QPDs to assess their flexibility and behavior. This comparative analysis can guide the selection of an appropriate distribution to characterize the quantity of interest. In Section 4, we explore several methods for extending univariate distributions to a

multivariate setting. These methods include the utilization of standard multivariate distributions, copulas, and bivariate quantiles. We show how these techniques can be applied to develop a bivariate version of the Generalized Myerson distribution and demonstrate its application in parametric and predictive elicitation. Finally, in Section 5, we discuss future research directions and potential applications of QPDs in Bayesian analysis.

## 2. Quantile Parameterization of Probability Distributions

A fundamental principle in Bayesian data analysis is that learning from data involves more than formulating hypotheses and models. It necessitates articulating prior beliefs, expressing existing knowledge mathematically, and translating it into probability distributions for model parameters.

To accurately translate knowledge into the language of statistical models, the encoding distribution needs to be flexible, the process should be transparent, and the results must be interpretable. For continuous distributions, elicitation often consists of capturing a series of quantile-probability pairs (QPPs) (Kadane and Wolfson 1998, Morgan 2014) and then fitting a distribution to these pairs (O'Hagan 2019). However, in practice, the choice of a parametric distribution to fit the elicited QPPs is often influenced by concerns about conjugacy with the selected statistical model that represents the data-generative process (the likelihood function) and/or the availability of required distribution functions and fitting algorithms in the software employed. Frequently, the selected distribution possesses fewer parameters than the number of elicited QPPs, which can result in a less-than-perfect fit (O'Hagan 2019). For instance, it is common to elicit three quantiles (the median along with an upper and lower quartile) and subsequently attempt to fit a normal or lognormal distribution (which features two parameters) to these points.

An alternative approach to characterizing the distribution of predictions or parameters is through quantile-parameterized distributions (QPDs). These distributions are parameterized by the QPPs, allowing the elicited values to directly define the distribution, thereby ensuring a good fit and interpretability of parameters. The QPDs examined in this paper can accommodate a wide

range of shapes and boundedness, making them valuable for accurately representing experts' prior beliefs.

QPDs are constructed using the *quantile function*, either by transforming simpler quantile functions or by simultaneous fitting of parameterizing quantiles, as described below.

Let  $Y$  be a random variable with a (cumulative) distribution function (CDF) denoted as  $F_Y(y | \theta)$ . The quantile function (QF)  $Q_Y(u | \theta)$  for  $Y$  is defined as

$$Q_Y(u | \theta) = \inf\{y : F_Y(y | \theta) \geq u\}, u \in [0, 1]$$

Here,  $\theta$  represents the distribution parameter, and the subscript  $Y$  indicates that the depth  $u$  corresponds to the observation of the random variable  $Y$ . The intermediate cumulative probability  $u$  is referred to as the *depth* because it indicates how “deep” an observation is into the distribution (Perepolkin et al. 2023).


Both the CDF and the QF are considered equally valid ways of defining a distribution (Tukey 1965). For a distribution function that is right-continuous and strictly increasing over the support of  $Y$ , the quantile function  $Q_Y(u)$  is simply the inverse of the distribution function, denoted as  $Q_Y(u | \theta) = F_Y^{-1}(u | \theta)$ . Therefore, the quantile function is often referred to as the *inverse CDF*.

The derivative of the quantile function, known as the *quantile density function* (QDF), is denoted as  $q(u) = \frac{dQ(u)}{du}$ . It is reciprocally related to the probability density function (PDF)  $f(x)$  such that  $f[Q(u)]q(u) = 1$ . The quantity  $f_Y[Q_Y(u | \theta)] = [q_Y(u | \theta)]^{-1}$  is referred to as the *density quantile function* (Parzen 1979) or *p-pdf* (Gilchrist 2000). The relationships between these functions are concisely illustrated in the probability function Möbius strip (Figure 1).

Although many of the distributions discussed in Section 3 have closed-form cumulative distribution functions (CDFs) and probability density functions (PDFs), the functional form of the quantile function (QF) is often simpler and can be reasoned about in terms of other quantile functions, following *Gilchrist's QF transformation rules* summarized in Table 1.

The quantile-parameterized distributions in this paper are categorized into two groups based on their construction method. The first group comprises distributions that are *directly* parameterized by the quantile-probability pairs (QPPs). This group includes the Myerson

**Figure 1.** (Color online) Möbius Strip of Probability Functions (Perepolkin et al. 2023)



distribution (Myerson 2005), and the Johnson Quantile-Parameterized Distribution (Hadlock and Bickel 2017, 2019). These distributions are constructed by reparameterizing or transforming existing distributions, following Gilchrist rules (Table 1). The transformations used to construct them are detailed in the next section.

The other group of distributions is *indirectly* parameterized by the QPPs. They require a fitting step where the quantile-probability pairs are translated into distribution parameters, usually through optimization or least-squares methods. This group includes the Simple Q-Normal (Keelin and Powley 2011), Metalog (Keelin 2016), and the quantile-parameterized Triangular (Two-Sided Power) distribution by Kotz and Van Dorp (2004). Each distribution's fitting method is described in the respective subsections below.

### 3. Univariate Quantile-Parameterized Distributions

This section reviews various continuous univariate QPDs and discusses the generalized form for these distributions based on the variations of these QPDs appearing in the literature. For each distribution, we present its quantile function and discuss the parameterization and feasibility conditions. The derivative and inverse of each distribution can be found in the Supplementary Materials.

**Table 1.** Gilchrist's Quantile Function Transformation Rules (Gilchrist 2000)

Original QF	Rule	Resulting QF	Resulting variable
$Q_Y(u)$	Reflection rule	$-Q(1-u)$	QF of $-Y$
$Q_Y(u)$	Reciprocal rule	$1/Q(1-u)$	QF of $1/Y$
$Q_1(u), Q_2(u)$	Addition rule	$Q_1(u) + Q_2(u)$	valid QF
$Q_1(u), Q_2(u)$	Linear combination rule	$aQ_1(u) + bQ_2(u)$	valid QF for $a, b > 0$
$Q_1(u), Q_2(u) > 0$	Multiplication rule	$Q_1(u)Q_2(u)$	valid QF
$Q_Y(u)$	Q-transformation	$T(Q_Y(u))$	QF of $T(Y)$ , $T(Y)$ nondecreasing
$Q_Y(u)$	p-transformation	$Q_Y(H(u))$	p-transformation of $Q_Y(u)$ , $H(u)$ nondecreasing

### 3.1. Myerson Distribution

One of the earliest examples of a distribution parameterized by quantiles is the *generalized log-normal* distribution defined by the median and the upper and lower quartiles proposed by Myerson (2005). It relies on a transformation of the normal quantile function.

The Myerson distribution can be viewed as parameterized by three quantile values  $\{q_1, q_2, q_3\}$ , which correspond to the cumulative probabilities  $\{\alpha, 0.5, 1 - \alpha\}$ . These quantiles are symmetrical around the median and are defined by the tail parameter  $0 < \alpha < 0.5$ . This type of parameterization is known as the Symmetric Percentile Triplet (SPT,  $\alpha$ -level SPT or  $\alpha$ -SPT) parameterization. It is also used in several other QPDs that we describe below. The Myerson quantile function is

$$\rho = q_3 - q_2; \beta = \frac{\rho}{q_2 - q_1}; \kappa(u) = \frac{S(u)}{S(1-\alpha)}$$

$$Q_Y(u | q_1, q_2, q_3, \alpha) = \begin{cases} q_2 + \rho \frac{\beta^{\kappa(u)} - 1}{\beta - 1}, & \beta \neq 1 \\ q_2 + \rho \kappa(u), & \beta = 1 \end{cases}$$

Here,  $u$  represents the depth of the observations of the random variable  $Y$  given the parameterizing  $\alpha$ -SPT  $\{q_1, q_2, q_3, \alpha\}$ , with  $0 < \alpha < 0.5$ . The parameter  $\rho$  is the *upper p-difference*, and  $\beta$  is the ratio of the inter-percentile ranges, known as the *skewness ratio* (Gilchrist 2000, p. 72). The *kernel* quantile function  $S(u)$  is equal to the quantile function of the standard normal distribution, also known as the *probit*, defined as  $S(u) = \Phi^{-1}(u)$ , where  $\Phi(Y)$  denotes the CDF of the standard normal distribution. The formulas for the derivative and the inverse quantile function of the Myerson QPD can be found in the Supplementary Materials.

It is important to note that although the Myerson distribution includes the normal distribution as a

special case when the skewness parameter  $\beta = 1$ , it can exhibit right-skewness or left-skewness for other values of  $\beta$ . In the symmetrical case, the range of the quantile function is  $(-\infty, \infty)$ . For the right-skewed distribution ( $\beta > 1$ ), the range is  $(q_2 - \frac{\rho}{\beta-1}, \infty)$ , and for the left-skewed distribution ( $0 < \beta < 1$ ), the range is  $(-\infty, q_2 - \frac{\rho}{\beta-1})$ . The limiting case of the skewed Myerson distribution  $\lim_{u \rightarrow 0} Q_Y(u | \theta)$  for  $\beta > 1$  (and the other limit for  $0 < \beta < 1$ ) possesses some important properties that we discuss in Section 3.3.2 below.

The basic quantile function (Gilchrist 2000, Lampasi 2008)  $S(u)$  underlying the Myerson distribution is a simple *probit*,  $S(u) = \Phi^{-1}(u)$ , transformed using the exponentiation function  $T(x) = \beta^x$  (Table 1), where  $\beta > 0$  represents the skewness ratio (Gilchrist 2000). The quantile parameterization is facilitated by  $\kappa(u)$ , which takes values  $\{-1, 0, 1\}$  for the three quantiles  $\{q_1, q_2, q_3\}$  such that  $Q(\alpha) = q_1$ ,  $Q(0.5) = q_2$ , and  $Q(1 - \alpha) = q_3$ .

### 3.2. Johnson Quantile-Parameterized Distribution

Hadlock (2017) reviewed the existing quantile-parameterized distributions and proposed the quantile parameterization of the Johnson SU family of distributions (Johnson et al. 1994). Hadlock and Bickel (2017) presented two versions of the distribution: the bounded (J-QPD-B) and the semibounded (J-QPD-S), both parameterized by an SPT  $\{q_1, q_2, q_3, \alpha\}$  and the bound(s).

The J-QPD-B distribution is obtained by applying the inverse-probit transformation to the Johnson SU quantile function  $Q_{SU}(u) = \xi + \lambda \sinh[\delta(S(u) + \gamma)]$ , where  $\delta$  and  $\gamma$  are two shape parameters. This function is then rescaled to the compact interval  $[l_b, u_b]$ . The J-QPD-B quantile function is

$$Q_B(u) = \begin{cases} l + rS^{-1}[Q_{SU}(u)], & n \neq 0 \\ l + rS^{-1}[B + \kappa S(u)], & n = 0 \end{cases}$$

where

$$\begin{aligned}
 Q_{SU}(u) &= \xi + \lambda \sinh[\delta(S(u) + \gamma)] \\
 r &= (u_b - l_b); \quad \gamma = nc; \quad \kappa = \frac{H-L}{2c} \\
 S(u) &= \Phi^{-1}(u); \quad c = S(1 - \alpha); \\
 L &= S\left(\frac{q_1 - l_b}{u_b - l_b}\right); \quad B = S\left(\frac{q_2 - l_b}{u_b - l_b}\right); \\
 H &= S\left(\frac{q_3 - l_b}{u_b - l_b}\right); \quad n = \text{sgn}(L + H - 2B) \\
 \xi &= \begin{cases} L, & n = 1, \\ B, & n = 0, \\ H, & n = -1, \end{cases} \\
 \delta &= \frac{1}{c} \cosh^{-1}\left(\frac{H-L}{2\min(B-L, H-B)}\right) \\
 \lambda &= \frac{H-L}{\sinh(2\delta c)}
 \end{aligned}$$

The left panel in Figure 2 showcases the J-QPD-B quantile function, which is parameterized using 0.25-SPT and 0.01-SPT assessments of the proportion of fruit infested with *Citripestis sagittiferella*, as elicited by EFSA et al. (2023). The dashed line represents the Beta distribution fitted by the authors. The J-QPD-B, being parameterized by an SPT, effectively captures three of the five parameterizing quantiles, whereas the Beta distribution provides only an approximation. Besides, finding parameters of the Beta distribution requires an optimization step.

The J-QPD-S distribution is a semi-bounded variant of the distribution that employs exponentiated hyperbolic arcsine transformations of the Johnson’s SU

quantile function  $Q_{Sua}(u) = \xi + \lambda \sinh[\text{asinh}(\delta S(u)) + \text{asinh}(\delta\gamma)]$  located at  $\xi = 0$  (Hadlock and Bickel 2017)

$$Q_S(u) = \begin{cases} l_b + \theta \exp[Q_{Sua}(u)], & n \neq 0 \\ l_b + \theta \exp[\lambda \delta S(u)], & n = 0 \end{cases}$$


where

$$\begin{aligned}
 Q_{Sua}(u) &= \lambda \sinh[\text{asinh}(\delta S(u)) + \text{asinh}(\delta\gamma)] \\
 r &= (u_b - l_b); \quad \gamma = nc; \quad \kappa = \frac{H-L}{2c} \\
 S(u) &= \Phi^{-1}(u); \quad c = S(1 - \alpha); \\
 L &= \ln(q_1 - l_b); \quad B = \ln(q_2 - l_b); \\
 H &= \ln(q_3 - l_b); \quad n = \text{sgn}(L + H - 2B) \\
 \theta &= \begin{cases} q_1 - l_b, & n = 1, \\ q_2 - l_b, & n = 0, \\ q_3 - l_b, & n = -1, \end{cases} \\
 \delta &= \frac{1}{c} \sinh\left[\cosh^{-1}\left(\frac{H-L}{2\min(B-L, H-B)}\right)\right] \\
 \lambda &= \frac{1}{\delta c} \min(H-B, B-L)
 \end{aligned}$$

When  $n = \text{sgn}(L + H - 2B)$  evaluates to zero, the result is a lognormal distribution with parameters  $\mu = \ln(\theta) = \ln(q_2 - l_b)$  and  $\sigma = \lambda \delta = (H - B)/c$ . This distribution has support on the interval  $[l_b, \infty]$ .

The right panel in Figure 2 depicts the J-QPD-S quantile function, which is parameterized using 0.25-SPT and 0.01-SPT assessments of the total trade flow for citrus fruit imported by the EU from Indonesia, Malaysia, Thailand, and Vietnam in tons/year (EFSA et al. 2023).

**Figure 2.** (Color online) Fitted J-QPD-B (Left) and J-QPD-S (Right) Distribution for Prevalence at Origin and Total Trade Flow, Respectively



### 3.3. Generalizations of QPDs

**3.3.1. Generalized Johnson Quantile-Parameterized Distribution.** Hadlock and Bickel (2019) introduced the *generalized* version of the Johnson Quantile-Parameterized Distribution system, denoted as G-QPD, by replacing  $S(u)$  the probit at the core of the Johnson SU quantile function with the quantile functions of the logistic and Cauchy distributions. The standard quantile function and distribution function of the logistic distribution are given by  $S(u) = \ln\left(\frac{u}{1-u}\right)$ ,  $F(y) = [\exp(-y) + 1]^{-1}$ . The standard quantile function and distribution function of the Cauchy distribution are given by  $S(u) = \tan\left[\pi\left(u - \frac{1}{2}\right)\right]$ ,  $F(y) = \frac{1}{\pi} \arctan(y) + \frac{1}{2}$ .

Hadlock and Bickel (2019) showed that the *kernel* quantile function  $S(u)$  can be any standardized ( $S(0.5) = 0$ ), symmetrical ( $s(u) = s(1-u)$ ), and unbounded ( $S(u) \in (-\infty; \infty)$ ) quantile function with a smooth quantile density  $dS(u)/du = s(u)$ . These authors further showed that if  $S(u)$  and  $S^{-1}(y)$  are expressible in closed-form, then the quantile function and distribution function of G-QPD will also be closed-form.

For the logistic kernel, the G-QPD-S represents the generalized log-logistic distribution, characterized by two shape parameters,  $\lambda$  and  $\delta$ . For the Cauchy kernel, the G-QPD-S corresponds to the shifted log-Cauchy distribution (Hadlock and Bickel 2019).

**3.3.2. Generalized Myerson Distributions.** Following the approach in Hadlock and Bickel (2019), the Myerson distribution can be generalized by substituting the normal kernel quantile function  $S(u) = \Phi^{-1}(u)$  with an

alternative symmetrical quantile function based on the depth  $u$ . Below, we discuss possible kernels and the resulting distributions:


**3.3.2.1. Logit-Myerson Distribution.** Recently, Wilson et al. (2023) reparameterized the log-logistic distribution in terms of a symmetric percentile triplet. Even though the authors did not recognize it as such, the resulting quantile-parameterized distribution was a Myerson distribution with the logit kernel QF  $S(u) = \ln\left(\frac{u}{1-u}\right)$ .

There could be several reasons why one might prefer the logit function over the probit function (Berkson 1951). For example, distribution based on logit may exhibit greater numerical stability because of its simple closed-form quantile function, which does not rely on numerical approximation during sampling. Logit-Myerson distribution displays slightly heavier tails compared with the standard (probit-based) Myerson distribution (Figure 3).

**3.3.2.2. Sech-Myerson Distribution.** Following the same principle, a variant of Myerson distribution may be created using the hyperbolic secant quantile function  $S(u) = \ln\left[\tan\left(\frac{\pi}{2}u\right)\right]$ .

The Sech-Myerson distribution possesses even thicker tails than the Logit-Myerson distribution for the same parameterizing SPT  $\{-5, 4, 16, 0.25\}$  (Figure 3). In Section 3.6, we conduct a comparative analysis of different variations of the Generalized Myerson distribution alongside their parametric counterparts and other quantile distributions.

**Figure 3.** (Color online) Quantile Function and Quantile Density of Generalized Myerson Distributions



Theoretically, there is an infinite range of quantile function kernels that can be utilized to generate new variations of the Generalized Myerson distribution. These candidate kernel distributions can even include shape parameters as long as the resulting  $S(u)$  remains standardized, symmetrical, and unbounded, as specified above. For instance, it is possible to incorporate the basic QF of the Tukey Lambda distribution  $S(u | \lambda) = u^\lambda - (1 - u)^\lambda$  for a fixed  $\lambda \neq 0$ , or the Cauchy distribution  $S(u) = \tan[\pi(u - 0.5)]$ , as employed by Hadlock and Bickel (2019). However, not all standard quantile functions are created equal. To illustrate the issue of unreliable kernels, let us consider Myerson distributions based on the Cauchy and Tukey Lambda quantile functions (for  $\lambda = -0.5$ ). As can be observed in Figure 4, the density of the Generalized Myerson distribution with these kernels exhibits unexpected spike near the lower bound.

Whereas all right-skewed Generalized Myerson distributions are bounded on the left at  $\lim_{u \rightarrow 0} Q(u | \theta) = q_2 - \rho \frac{1}{\beta - 1}$  regardless of the kernel used, the quantile density at the left limit  $\lim_{u \rightarrow 0} [q(u | \theta)]^{-1}$  is not independent of the kernel. Although we can assume that  $q(0) = \infty$ , the lower tail of the density quantile function  $[q(u)]^{-1}$  may exhibit a curling effect for certain kernels, resulting in an increase in density for lower values of  $u$ . This effect is caused by the nonmonotonic behavior of the quantile convexity function  $c(u) = dq(u)/du$ . This can be easily verified by taking the second derivative of  $\beta^{S(u)}$  for  $\beta > 0$ . Although such kernels are mathematically valid and yield a nondecreasing Generalized Myerson QF, we believe that they may be less useful

because of the counterintuitive concentration of density in the bounded tail. Consequently, we do not recommend using Cauchy or Tukey Lambda kernels in practical applications.

### 3.4. Simple Q-Normal, Metalog Distributions

An alternative system of quantile-parameterized distributions was proposed by Keelin and Powley (2011) and Powley (2013). Their approach relied on the finite Taylor expansion of parameters in standardized quantile functions. Within this framework, two distributions were introduced: the Simple Q-Normal distribution and the Metalog distribution.


The Simple Q-Normal (SQN) distribution was developed by expanding the parameters in the normal quantile function. Keelin and Powley (2011) used this method to express the parameters of the normal quantile function  $Q(u | \mu, \sigma) = \mu + \sigma z(u)$  as linear functions of the depth  $u$ . Specifically,  $\mu(u) = a_1 + a_4 u$  and  $\sigma(u) = a_2 + a_3 u$ , where  $z(u) = \Phi^{-1}(u)$  denotes the standard normal quantile function. Therefore, the quantile function of the SQN distribution can be expressed as

$$Q(u) = a_1 + a_2 z(u) + a_3 u z(u) + a_4 u$$

where  $z(u) = \Phi^{-1}(u)$ , and  $\mathbf{a} = \{a_1, a_2, a_3, a_4\}$  represents a vector of parameters.

Consider a quantile-probability tuple of size 4, denoted as  $\{\mathbf{p}, \mathbf{q}\}_4$ , which consists of an ordered vector of cumulative probabilities  $\mathbf{p} = \{p_1, p_2, p_3, p_4\}$  and an ordered vector of corresponding quantiles  $\mathbf{q} = \{q_1, q_2, q_3, q_4\}$  (Perepolkin et al. 2024). Substituting these vectors into the SQN quantile function for  $u$  and  $Q(u)$ ,

**Figure 4.** (Color online) Quantile Function and Quantile Density of Generalized Myerson Distributions with Unreliable Kernels



respectively, we obtain the matrix equation  $\mathbf{q} = \mathbb{P}\mathbf{a}$ , where  $\mathbf{a} = \{a_1, a_2, a_3, a_4\}$  represents the parameter vector of the SQN distribution and

$$\mathbb{P} = \begin{bmatrix} 1 & z(p_1) & p_1 z(p_1) & p_1 \\ 1 & z(p_2) & p_2 z(p_2) & p_2 \\ 1 & z(p_3) & p_3 z(p_3) & p_3 \\ 1 & z(p_4) & p_4 z(p_4) & p_4 \end{bmatrix}.$$

The parameter vector  $\mathbf{a}$  can be obtained by solving the matrix equation above, given the four-element quantile-probability tuple  $\{\mathbf{p}, \mathbf{q}\}_4$  (Keelin and Powley 2011, Perepolkin et al. 2024).

The same approach was later employed by Keelin (2016) in creating the metalog (meta-logistic) distribution. Starting with the quantile function of the logistic distribution  $Q(u | \mu, s) = \mu + \text{slogit}(u)$ , where  $\mu$  corresponds to the mean and  $s$  is proportional to the standard deviation  $\sigma = s\pi/\sqrt{3}$ , Keelin (2016) expanded the parameters  $\mu$  and  $s$  using a finite Taylor series centered at 0.5. Specifically,  $\mu(u) = a_1 + a_4\check{u} + a_5\check{u}^2 + \dots$ , and  $s(u) = a_2 + a_3\check{u} + a_6\check{u}^2 + \dots$ , where  $\check{u} = u - 0.5$  and  $a_i, i = \{1, 2, \dots, n\}$  are real constants.

Therefore, the metalog quantile function is

$$Q(u) = a_1 + a_2 \text{logit}(u) + a_3 \check{u} \text{logit}(u) + a_4 \check{u} + a_5 \check{u}^2 \dots,$$

where  $\check{u}$  is the centered depth  $\check{u} = u - 0.5$ , and  $\text{logit}(u) = \ln(u) - \ln(1 - u)$ .

Given a QPT of size  $m$  denoted by  $\{\mathbf{p}, \mathbf{q}\}_m$ , where  $\mathbf{p}$  and  $\mathbf{q}$  are ordered vectors of cumulative probabilities and corresponding quantiles, respectively, the vector of coefficients  $\mathbf{a} = a_1, \dots, a_m$  can be determined by solving the matrix equation  $\mathbf{q} = \mathbb{P}\mathbf{a}$ , where  $\mathbf{p}$ ,  $\mathbf{q}$ , and  $\mathbf{a}$  are column vectors, and  $\mathbb{P}$  is an  $m \times n$  matrix:

$$\mathbb{P} = \begin{bmatrix} 1 & \text{logit}(p_1) & \check{p}_1 \text{logit}(p_1) & \check{p}_1 & \dots \\ 1 & \text{logit}(p_2) & \check{p}_2 \text{logit}(p_2) & \check{p}_2 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & \text{logit}(p_m) & \check{p}_m \text{logit}(p_m) & \check{p}_m & \dots \end{bmatrix}.$$

The vector of coefficients  $\mathbf{a}$  can be determined as  $\mathbf{a} = [\mathbb{P}^T \mathbb{P}]^{-1} \mathbb{P}^T \mathbf{q}$ . If  $\mathbb{P}$  is a square matrix, meaning that the number of terms  $n$  is equal to the size of the parameterizing QPT  $m$ , then the equation can be further simplified to  $\mathbf{a} = \mathbb{P}^{-1} \mathbf{q}$ . Metalog is said to be *approximated* when the number of quantile-probability pairs used for parameterization exceeds the number of terms in the metalog QF (Keelin 2016, Perepolkin et al. 2024).

The SQN and Metalog distributions are families of extended distributions that, in theory, can have an arbitrary number of terms. Keelin (2016) demonstrated the flexibility of the metalog distribution and its ability to approximate arbitrarily complex probability density functions with high precision, given enough terms in the metalog specification. In practice, 10 – 15 terms are sufficient to approximate the distributional shapes of virtually any complexity (Keelin and Howard 2021). Keelin (2016) introduced the bounded logit-metalog, the semi-bounded log-metalog, and a special case of a three-term metalog parameterized by  $\alpha$ -SPT (SPT-metalog).

However, not all combinations of parameters  $\mathbf{a}$  in metalog and SQN distributions result in a feasible (nondecreasing) quantile function. For an arbitrary  $\mathbf{a}$ -vector, feasibility must be checked (Keelin and Powley 2011). In the case of three-term metalogs, the feasibility conditions are straightforward (Keelin 2016). However, as the number of terms increases, these conditions become increasingly complex (Keelin 2017). Dealing with such feasibility requirements stands in contrast with QFs that are constructed using Gilchrist rules (Table 1), which guarantee feasibility.

### 3.5. Other Distributions

Many classical parametric distributions can be identified using a set of quantiles, although they may not be strictly “quantile-parameterized.” That is, these distributions might not have a closed-form quantile function expressed in terms of a Quantile-Probability Tuple (QPT), but they can still have a unique set of parameters corresponding to a QPT of matching size.

Cook (2010) demonstrated that any distribution from the location-scale family can be parameterized using two quantiles. Consider a distribution with a quantile function  $Q(u) = \mu + \sigma S(u)$ , where  $S(\cdot)$  is a basic quantile function. Given a quantile-probability tuple  $\{\mathbf{p}, \mathbf{q}\}_2$  composed of quantile values  $\mathbf{q} = \{q_1, q_2\}$  and probabilities  $\mathbf{p} = \{p_1, p_2\}$ , where  $x_1 < x_2$  and  $p_1 < p_2$ , the scale parameter  $\sigma$  can be computed as

$$\sigma = \frac{q_2 - q_1}{S(p_2) - S(p_1)}$$

Similarly, the location parameter  $\mu$  can be computed as

$$\mu = \frac{q_1 S(p_2) - q_2 S(p_1)}{S(p_2) - S(p_1)}$$

This means that distributions from the location-scale family can be expressed using the quantile function

$$Q(u) = \frac{[q_1 S(p_2) - q_2 S(p_1)] + [q_2 - q_1] S(u)}{S(p_2) - S(p_1)}$$

Cook (2010) also provides examples of quantile-matching for other two-parameter distributions, where one or both parameters may be shape parameters. Some distributions, like the Weibull, have closed-form solutions, whereas others, such as the Beta distribution, require numerical optimization or root-finding algorithms. Van Dorp and Mazzuchi (2000) proved the existence of a solution for the Beta distribution parameters based on two quantiles, whereas Shih (2015) demonstrated the uniqueness of that solution. Van Dorp and Mazzuchi (2000) outlined the procedure for determining Beta distribution parameters using two quantiles.

Kotz and Van Dorp (2004) described the procedure for quantile parameterization of the triangular distribution (Johnson 1997). This bounded distribution is widely used in finance and insurance and has been popularized by the @Risk software package (Palisade Corp 2009). The triangular distribution is parameterized by the two quantiles  $q_a$  and  $q_b$  and the mode  $m$ , subject to the constraint that  $a \leq q_a \leq m \leq q_b \leq b$ , where  $a$  and  $b$  represent the lower and upper bounds, respectively. The standard quantile function for the triangular distribution is expressed in terms of the bounds  $a$  and  $b$  and the mode  $m$ ,

$$Q(u) = \begin{cases} a + \sqrt{ud_1 d}, & \text{for } 0 \leq u \leq \frac{d_1}{d} \\ b - \sqrt{(1-u)d_u d}, & \text{otherwise,} \end{cases}$$

where  $d = b - a$ ;  $d_1 = m - a$ ;  $d_u = b - m$ .

Kotz and Van Dorp (2004) showed that given the two parameterizing quantile-probability pairs  $\{q_a, p_a\}$  and  $\{q_b, p_b\}$  and the mode value  $m$ , there exists a unique value of depth  $p_a < p < p_b$  corresponding to the root of the function

$$g(p) = \frac{(m - q_a)(1 - \tau_b)}{(q_b - m)(1 - \tau_a) + (m - q_a)(1 - \tau_b)} - p$$

where  $\tau_b = \sqrt{\frac{1-p_b}{1-p}}$ ,  $\tau_a = \sqrt{\frac{p_a}{p}}$ .

The root value  $p \in (p_a, p_b)$  of the function  $g(p)$  can be found using any of the bracketing root-finding algorithms (Perepolkin et al. 2023). It can then be substituted into the following expressions to find the lower  $a$  and

upper  $b$  limit parameters of the triangular distribution

$$a(p) \equiv \frac{q_a - m\tau_a}{1 - \tau_a}, \quad a(p) < q_a$$

$$b(p) \equiv \frac{q_b - m\tau_b}{1 - \tau_b}, \quad b(p) > q_b$$

where  $\tau_b = \sqrt{\frac{1-p_b}{1-p}}$ , and  $\tau_a = \sqrt{\frac{p_a}{p}}$ .

Section 4.3.3 in Kotz and Van Dorp (2004) provides an algorithm for fitting a four-parameter generalization of the triangular distribution, called the Two-Sided Power Distribution (TSP), using three quantile-probability pairs and a mode value. By allowing the TSP power parameter  $n \rightarrow \infty$ , Kotz and van Dorp (2005) developed an algorithm for identifying the parameters of the Asymmetric Laplace distribution with unbounded support. Additionally, van Dorp and Mazzuchi (2021) proved that a lower quantile, a mode, and an upper quantile uniquely determine the left and right branch power parameters  $m, n > 0$  of a Generalized Two-Sided Power (GTSP) distribution (Herrerías-Velasco et al. 2009), given a fixed bounded support  $[a, b]$ . An algorithm to solve for the power parameters  $m, n > 0$  of the GTSP distribution is also provided in van Dorp and Mazzuchi (2021). More recently, van Dorp and Shittu (2023) applied the same algorithm to solve for the left and right branch power parameters  $m, n > 0$  of the Generalized Two-Sided Beta (GTSB) distribution, which serves as a smooth alternative to the GTSP distribution.

### 3.6. Choosing Quantile-Parameterized Distribution

A common approach to assess the properties of probability distributions is through central moments, denoted by  $\mu_k = \mathbb{E}[(Y - \mu)^k]$ , where  $\mu$  represents the expected value of  $Y$ . Karl Pearson introduced a classification system for distributions using moment ratios associated with skewness and kurtosis (Fiori and Zenga 2009):

$$\beta_1 = \frac{\mu_3^2}{\mu_2^3}, \quad \beta_2 = \frac{\mu_4}{\mu_2^2}$$

Although computing moments using the quantile function is straightforward (the  $n$ -th raw moment is  $\mu_k = \int_0^1 Q(u)^k du$ ), it may not be possible to calculate higher-order moments for certain distributions.

Instead, robust alternatives to moments can be utilized, such as the sample median  $\mu_r$ , the interquartile range  $\sigma_r$ , the quartile-based robust coefficient of skewness  $s_r$  (Kim and White 2004), also known as Bowley's

skewness (Bowley 1920) or Galton’s skewness (Gilchrist 2000), and the octile-based robust coefficient of kurtosis  $\kappa_r$ , also known as Moors’ kurtosis (Moors 1988):

$$\begin{aligned} \mu_r &= Q(1/2) \\ \sigma_r &= Q(3/4) - Q(1/4) \\ s_r &= \frac{Q(3/4) + Q(1/4) - 2Q(1/2)}{\sigma_r} \\ \kappa_r &= \frac{Q(7/8) - Q(5/8) + Q(3/8) - Q(1/8)}{\sigma_r} \end{aligned}$$

Kim and White (2004) and Arachchige et al. (2022) have proposed to standardize robust moments to facilitate their comparison with the corresponding robust moments of the standard normal distribution. Groeneveld (1998) and Jones et al. (2011) have introduced generalizations of robust moments to other quantiles.


Unlike moments, quantiles are always well defined, and because QPDs are parameterized by quantile-probability pairs, quantile-based robust moments can sometimes be directly computed from the parameters. For instance, if the basic quantile function  $S(u)$  in  $Q(u) = \mu + \sigma S(u)$  is standardized (such that  $S(0.5) = 0$ ), where  $\mu$  and  $\sigma$  are the location and scale parameters, respectively, of  $Q(u)$  then  $\mu_r = \mu$ . Moreover,  $\sigma_r$  is always independent of location, and  $s_r$  and  $\kappa_r$  are independent of both location and scale.

Figures 5, 6, and 7 resemble the Cullen et al. (1999) plots (also known as the Pearson plots), but instead of using central moments they employ quartile/octile-based robust metrics of skewness  $s_r$  and kurtosis  $\kappa_r$  to compare the quantile-parameterized distributions to some of their parametric counterparts.


In these plots, Metalog3 and Metalog4 refer to three- and four-term metalog distributions, respectively, and GLDcsw refers to Chalabi et al. (2012) parameterization of the Generalized Lambda Distribution (GLD). As can be seen in Figure 5, all generalizations of Myerson distributions have higher robust kurtosis for the same robust skewness, which indicates heavier tails outside of parameterizing quantiles. Additionally, GLD CSW is more flexible than the unbounded four-term metalog. The *log*-transformed metalog distribution appears to be the most flexible among the semi-bounded distributions (Figure 6). Furthermore, the flexibility of the bounded J-QPD-B is at least as good as that of the Beta and Kumaraswamy distributions (Figure 7).

The practical choice of a distribution is often guided by information elicited from an expert. Many elicitation protocols result in a symmetric triplet of quantile-probability pairs (SPT), such as the 25th, 50th, and 75th percentiles or the 10th, 50th, and 90th percentiles. In

**Figure 5.** (Color online) Robust Skewness vs. Robust Kurtosis for Some Unbounded Distributions



**Figure 6.** (Color online) Robust Skewness vs. Robust Kurtosis for Some Left-Bounded Distributions





such cases, a wide variety of QPDs are available to encode quantities of interest based on different boundedness conditions.

Figure 8 illustrates the selection of SPT-parameterized distributions for each type of boundedness. Note that

whereas Myerson distributions have implicit lower and upper bounds, the other SPT-parameterized distributions have explicit bounds, which must be provided by the expert. The distributions are listed roughly in increasing order of robust kurtosis for the same level of robust

**Figure 7.** (Color online) Robust Skewness vs. Robust Kurtosis for Some Bounded Distributions



**Figure 8.** (Color online) Concept Map for Choosing Quantile-Parameterized Distribution

skewness. This consideration may prove useful during the validation phase of the elicitation process, especially when the assessor wishes to query the expert about a higher or lower quantile outside the initial parameterizing triplet. If the tail of the originally selected distribution is found to be too light (i.e., the expert provides a higher or lower quantile that exceeds the value implied by the selected distribution), then the next distribution in the ordering of robust kurtosis might be more appropriate. Note that whereas this additional quantile may not be matched exactly, the newly chosen distribution might approximate it more closely.

If the elicited quantiles are asymmetrical around the median (Figure 8), or if a higher number of quantiles need to be matched (e.g., the assessor decides to match an additional upper or lower quantile along with the previously elicited SPT), then the Metalog distribution is the only available option (out of the QPDs with an explicit quantile function discussed in this paper). This highly flexible distribution supports all types of distribution boundedness and can accommodate more than a dozen quantile-probability pairs. However, a limitation of the

Metalog distribution is that it may not always be feasible for any given QPT, meaning that in practice some quantiles might need to be approximated rather than matched exactly.

All distributions presented in this paper are implemented in qpd package (Perepolkin 2019) in R. In addition, Metalog implementation can be found in rmetalog (Faber and Jung 2021) and pymetalog (Smith et al. 2021) for R and Python, respectively. J-QPD distributions are available through rjqpd package (Ingram 2020) in R and jqpd package (Khan 2020) in Python.

#### 4. Multivariate Quantile-Parameterized Distributions

Quantile-parameterized distributions can serve as marginal distributions in multivariate models, where the dependency structure is captured by a standard (parametric) multivariate distribution, a copula, or described by bivariate quantiles. The marginal distributions alone are insufficient to determine the corresponding bivariate distribution, resulting in an infinite number of bivariate distributions with the same

margins (Gumbel 1960, 1961). In this section, we describe several methods for extending the distributions parameterized by the quantile-probability pairs to become Multivariate Quantile-Parameterized Distributions (MQPDs).

#### 4.1. MQPDs Based on Standard Multivariate Distributions

**4.1.1. Normal Distribution.** In the simplest case, multivariate Quantile-Parameterized Distributions (MQPDs) can be created using the multivariate normal distribution, following the approach of Hoff (2007). The Myerson, J-QPD, and SQN quantile functions are Q-transformations of the probit  $Q(z(u) | \theta)$ , where  $z(u) = \Phi^{-1}(u)$  represents the standard normal quantile function. The multivariate versions of these distributions can be viewed as the Q-transformations of the multivariate normal distribution. To extend these QPDs to  $J$  dimensions using the multivariate normal distribution, we employ the method outlined in Drovandi and Pettitt (2011).

The  $i$ -th component of a single observation  $y_i$  can be described by the quantile function  $y_i = Q(z(u_i) | \theta_i)$ , for  $i = 1, \dots, J$ , where  $\theta_i$  represents the set of parameters for component  $i$  (e.g.,  $\{q_1, q_2, q_3, \alpha\}_i$ ) for Myerson or J-QPD distributions). The vector  $[z(u_1), \dots, z(u_j)]^T \sim N(0, \Sigma)$ , where  $\Sigma$  denotes the covariance matrix. For invertible distributions, the inverse quantile function is the cumulative distribution function (CDF)  $Q^{-1}(y_i | \theta) = F(y_i | \theta)$ ; otherwise, the inverse can be computed numerically as  $\widehat{F}(y_i | \theta) = \widehat{Q}^{-1}(y_i | \theta)$  (Perepolkin et al. 2023).

Drovandi and Pettitt (2011) showed that the joint density of a single (multivariate) observation  $(y_1, \dots, y_J)$  can be expressed as

$$f(y_1, \dots, y_J | \theta) = \varphi[z(Q^{-1}(y_1 | \theta_1)), \dots, z(Q^{-1}(y_J | \theta_J)); \Sigma] \times \prod_{i=1}^J \frac{dQ^{-1}(y_i | \theta_i)}{dy_i}$$

where  $z(Q^{-1}(y_i | \theta_i)) = z_i$ ,  $\varphi(z_1, \dots, z_J; \Sigma)$  represents the multivariate normal density with a mean of zero and a covariance matrix of  $\Sigma$ , and  $\frac{dQ^{-1}(y_i)}{dy_i} = f(y_i)$  is the probability density function (PDF) of the QPD (refer to the Supplementary Materials).

For distributions without a PDF, the same joint density can be expressed as a joint *density quantile function*,

$$[q(u_1, \dots, u_j)]^{-1} = \varphi[z(u_1), \dots, z(u_j); \Sigma] \times \prod_{i=1}^J [q(u_i | \theta_i)]^{-1},$$

because  $Q^{-1}(y_i | \theta_i) = u_i$  and  $f(y_i | \theta_i) = [q(u_i | \theta_i)]^{-1}$  (Gilchrist 2000).

It is worth noting that this method of creating multivariate distributions does not require every component to follow the same distributional form. As illustrated earlier, it is entirely possible to combine several different QPDs using the multivariate Gaussian distribution (Drovandi and Pettitt 2011).

To use the MQPD for the prior, both the density of the multivariate normal and the marginal densities need to be explicitly added to the log-likelihood. This is possible when the marginal QPDs used to define the multivariate prior are invertible, such as Myerson and J-QPD, because both the CDF  $Q^{-1}(y_i | \theta_i)$  and PDF  $dQ^{-1}(y_i | \theta_i)/dy_i$  are required. When a quantile-based prior specification is used, only the multivariate normal log-density needs to be added because the Jacobian for the marginal QF transformation is reciprocal to the DQF of the prior (Perepolkin et al. 2023).

**4.1.2. Logistic Distribution.** The same approach of joining the marginal QPDs can be applied by using the base quantile functions of other distributions. For instance, the Logit-Myerson distribution (Wilson et al. 2023) is based on the logistic quantile function. Two Logit-Myerson distributions can be connected using the bivariate logistic distribution. Gumbel (1961) proposed three different formulations for the bivariate logistic distribution. The Type II distribution from the Morgenstern family (Sajeevkumar and Irshad 2014, Basikhasteh et al. 2021) has the following joint distribution and density functions:

$$\begin{aligned} F(y_1, y_2 | \beta) &= F_1(y_1)F_2(y_2) \\ &\quad \times [1 + \beta(1 - F_1(y_1))(1 - F_2(y_2))] \\ f(y_1, y_2 | \beta) &= f_1(y_1)f_2(y_2) \\ &\quad \times [1 + \beta(1 - 2F_1(y_1))(1 - 2F_2(y_2))] \end{aligned}$$

where  $F_i(y_i)$  and  $f_i(y_i)$  for  $i \in \{1, 2\}$  refer to the univariate logistic distribution and density functions, respectively, and  $-1 \leq \beta \leq 1$ . Because  $y_i = Q_i(u_i)$ , we can express the

bivariate density in the quantile form

$$\begin{aligned} f[Q(u_1), Q(u_2) | \beta] &= f_1[Q(u_1)]f_2[Q(u_2)] \\ &\quad \times [1 + \beta(1 - 2F_1[Q_1(u_1)])] \\ &\quad \times (1 - 2F_2[Q_2(u_2)])] \\ [q(u_1, u_2 | \beta)]^{-1} &= [q_1(u_1)]^{-1}[q_2(u_2)]^{-1} \\ &\quad \times [1 + \beta(1 - 2u_1)(1 - 2u_2)] \end{aligned}$$

For logistic distribution with  $Q(u) = \ln(u) - \ln(1 - u)$  and  $[q(u)]^{-1} = u(1 - u)$ , the bivariate logistic density quantile function can be expressed as

$$\begin{aligned} [q_L(u_1, u_2 | \beta)]^{-1} &= u_1(1 - u_1)u_2(1 - u_2) \\ &\quad \times [1 + \beta(1 - 2u_1)(1 - 2u_2)] \end{aligned}$$


If we combine the QPD marginals, the result is the joint quantile-based density for the bivariate logistic-based QPD, where the dependency is captured by the bivariate logistic distribution with the coupling parameter  $\beta$ , and the margins are QPDs. The joint density quantile function is given by

$$\begin{aligned} [q_{MQPD}(u_1, u_2 | \theta_1, \theta_2, \beta)]^{-1} &= u_1(1 - u_1)u_2(1 - u_2) \\ &\quad \times [1 + \beta(1 - 2u_1)(1 - 2u_2)] \\ &\quad \times [q_1(u_1 | \theta_1)]^{-1}[q_2(u_2 | \theta_2)]^{-1}, \end{aligned}$$

where  $[q_i(u_i | \theta_i)]^{-1}$ , for  $i = 1, 2$ , are the marginal QPD density quantile functions, such as the density quantile function (DQF) of the Logit-Myerson distribution (see the Supplementary Materials).

This way of constructing a bivariate distribution is illustrated in Figure 9 with the Bivariate Logit-Myerson Distribution, parameterized by  $\Theta = \{\theta_1, \theta_2, \rho\}$ , where the

**Figure 9.** (Color online) Density of Generalized Myerson Distributions Joined by Type II Bivariate Logistic Distribution



marginal Myerson distributions are given by  $y_{ij} = Q_j(z(u_{ij}), \theta_j)$  for  $j = 1, 2$ , with parameter vectors  $\theta_1 = \{3, 7, 10; 0.25\}$ ,  $\theta_2 = \{1, 10, 20; 0.1\}$ , and the dependence parameter  $\beta = 0.6$ .

#### 4.2. Copula-Based MQPDs

The approach we have used so far is similar to constructing the joint distribution using the Gaussian copula (Hoff 2007). Copulas provide a general approach to modeling joint distributions, separating the bivariate dependence from the effects of marginal distributions (Kurowicka and Cooke 2006). The literature describes a wide range of copulas (Genest and Favre 2007, Kurowicka and Joe 2011, Smith 2013), and new copulas can be created using generator functions (Durrleman et al. 2000). When a copula is used to connect QPDs, the joint density is calculated as

$$\begin{aligned} f_{MQPD}(y_1, y_2 | \theta_1, \theta_2, \Xi) &= c[F(y_1 | \theta_1), F(y_2 | \theta_2) | \Xi] \\ &\quad \times f_1(y_1 | \theta_1)f_2(y_2 | \theta_2), \end{aligned}$$

where  $c[\cdot]$  represents the copula density function with parameter  $\Xi$ , and  $F(y_i | \theta_i)$  and  $f_i(y_i | \theta_i)$  are the CDF and PDF, respectively, of the marginal quantile-parameterized distributions.


The same density can be expressed in a quantile-based form (Perepolkin et al. 2023):

$$\begin{aligned} [q_{MQPD}(u_1, u_2 | \theta, \Xi)]^{-1} &= c[u_1, u_2 | \Xi] \\ &\quad \times [q_1(u_1 | \theta_1)]^{-1}[q_2(u_2 | \theta)]^{-1}, \end{aligned}$$

where  $c[\cdot]$  is the copula density function with parameter  $\Xi$ , and  $[q_i(u_i | \theta_i)]^{-1}$ , for  $i = 1, 2$ , are the marginal DQFs of QPDs. Figure 10 presents 10,000 samples from the bivariate Myerson distribution joined by the Joe copula with  $\theta = 3$ .

Elicitation of multivariate distributions may require a specialized approach (Elfadaly and Garthwaite 2017, Wilson et al. 2021). For examples of expert-specified multivariate distributions encoded with copulas, we refer to the relevant literature (Aas et al. 2009, Sharma and Das 2018, Wilson 2018, Holzhauser et al. 2022). When fitting copulas to empirical observations, the *blanket* goodness of fit measure (Wang and Wells 2000) based on Kendall's transform (Genest et al. 2006, 2009) can be used.

**Figure 10.** (Color online) Samples from the Bivariate Myerson Distribution Joined by the Joe Copula ( $\theta = 3$ )



### 4.3. Bivariate Quantiles

The formal definition of bivariate quantile functions and the method for constructing bivariate quantile distributions using marginal and conditional quantile functions are provided by Nair and Vineshkumar (2023) and Vineshkumar and Nair (2019). They defined the bivariate quantile function (bQF) of  $(X_1, X_2)$  as the pair  $Q(u_1, u_2) = [Q_1(u_1), Q_{21}(u_2 | u_1)]$ , where  $Q_1(u_1) = \inf\{x_1 : F_1(x_1) \geq u_1\}$ ,  $u_1 \in [0, 1]$  and  $Q_{21}(u_2 | u_1) = \inf\{x_2 : F_{21}(Q_1, x_2) \geq u_2\}$ .

The conditional quantile function  $Q_{21}(u_2 | u_1)$  can be obtained by inverting the conditional distribution function  $F_{21}(x_1, x_2)$ , which is computed from the factorization of the joint survival function. The joint survival function is defined as  $\bar{F}(x_1, x_2) = P(X_1 > x_1)P(X_2 > x_2 | X_1 > x_1) = \bar{F}(x_1)\bar{F}_{21}(x_1, x_2)$ . Note that the joint survival function  $\bar{F}(x_1, x_2) = 1 - F_1(x_1) - F_2(x_2) + F(x_1, x_2)$ , and the conditional survival function  $\bar{F}_{21}(x_1, x_2) = 1 - F_{21}(x_1, x_2)$ .

Another approach for creating bivariate quantile functions is through Gilchrist’s QF transformation rules (Gilchrist 2000), which can be generalized to bivariate quantile functions. According to Nair and Vineshkumar (2023) (Property 6), the conditional QF can be constructed as a sum of two univariate QFs:  $Q_{21}(u_2 | u_1) = Q_1(u_1) + Q_2(u_2)$ . This means that the pair  $[Q_1(u_1); Q_1(u_1) + Q_2(u_2)]$  is a valid bivariate quantile function, which generalizes Gilchrist’s *addition rule* (Table 1). The addition rule also works for quantile density functions (Property 7). If  $Q_1$  is left-bounded at

zero, that is,  $Q_1(0) = 0$ , then the margins of such a bQF are  $X_1 = Q_1(u_1)$  and  $X_2 = Q_2(u_2)$ . Otherwise, the marginal distribution of  $X_2$  will be  $\lim_{u_1 \rightarrow 0} Q_{21}(u_2 | u_1)$ , which in many cases is not tractable.


If  $Q_1(u_1)$  and  $Q_2(u_2)$  are positive on  $u_i \in [0, 1]$ , then their product is also a valid conditional QF (Property 8), generalizing Gilchrist’s “product rule.” Finally, Property 9 generalizes the “Q-transformation rule,” stating that for every increasing transformation functions  $T_1$  and  $T_2$ ,  $[T_1(Q_1(u_1)), T_1(Q_1(u_1)) + T_2(Q_2(u_2))]$  is also a valid bQF.

Therefore, valid bivariate quantile-parameterized QFs can be created by constructing the conditional quantile functions as Gilchrist combinations of univariate quantile-parameterized QFs. Figure 11 shows 1,000 samples from the bivariate distribution created by adding together two Myerson distributions. Note that in this case, only the marginal distribution of  $x_1 = Q_1(u_1)$  is available in closed form:

$$\begin{aligned} (u_1, u_2) & \overset{X_1, X_2}{\sim} [Q_1(u_1), Q_1(u_1) + Q_2(u_2)] \\ Q_1(u_1) & \sim \text{Myerson}(3, 7, 10; 0.1) \\ Q_2(u_2) & \sim \text{Myerson}(-9, -3, 2; 0.25) \end{aligned}$$

This bQF is easy to elicit and interpret, because  $Q_2(u_2)$  can be thought of as a random adjustment to the value of  $Q_1(u_1)$ . In fact, the conditional quantile function  $Q_{21}(u_2 | u_1)$  can be thought of as having the classical form  $Q_{21}(u_2 | u_1) = \mu(u_1) + \sigma Q_2(u_2)$  (Gilchrist 2000), where the location is randomly varying with  $\mu(u_1) = Q_1(u_1)$  and the scale parameter  $\sigma = 1$ . First, the marginal distribution  $Q_1(u_1)$  is elicited, and then the

**Figure 11.** (Color online) Samples from the Bivariate Myerson Quantile Function



difference between the values  $x_1$  and  $x_2$  can be elicited as a QPT and encoded as  $Q_2(u_2)$ .

## 5. Discussion

Quantile-based distributions have garnered significant attention in the research community. Several distributions, such as the Generalized Lambda Distribution (GLD) (Ramberg and Schmeiser 1974, Freimer et al. 1988), the g-and-k distribution (Haynes et al. 1997, Haynes and Mengersen 2005, Prangle 2017), the g-and-h distribution (Mac Gillivray 1992, Rayner and MacGillivray 2002, Field and Genton 2006), and the Wakeby distribution (Jeong-Soo 2005, Tarsitano 2005, Rahman et al. 2015), have been studied extensively and documented in the literature. These distributions are defined by noninvertible quantile functions (Perepolkin et al. 2023). However, the research on quantile-parameterized distributions remains relatively unexplored. These distributions offer interpretable parameters that are defined on the same scale as the quantities of interest, simplifying the elicitation process for experts. Many popular elicitation protocols for both predictive and parametric elicitation rely on the assessment of quantile-probability pairs (QPPs). Instead of fitting a parametric distribution to the elicited QPPs (O'Hagan 2019, Best et al. 2020), assessors could directly use the elicited QPPs as inputs into one of the QPD quantile functions, which can be easily employed in both quantile-parameterized and parametric models.

Provided that the expert and the elicitor agree on the scientific model to be used for representing the expert's understanding of the world (Burgman et al. 2021), several types of inputs may be required to inform the model. Among those are the expert's judgement about the model *parameters* (O'Hagan 2019, Mikkola et al. 2023) and their *predictions* of the next observation (Winkler 1980, Kadane and Wolfson 1998, Akbarov 2009). Both parametric and predictive judgments should be captured together with corresponding uncertainties to reflect the expert's state of knowledge.

Quantile-parameterized distributions offer distinct advantages as high-fidelity priors that precisely capture expert assessments. These distributions are particularly beneficial for domain experts who may not be well-versed in statistics, because they provide high

flexibility while retaining parameter interpretability. As a result, QPDs can faithfully represent an expert's beliefs without compromising convenience or precision.

Quantile-parameterized distributions with unbounded support do not control the heaviness of the tail beyond the most extreme quantile specified. Therefore, different QPDs fitted to the same set of quantile-probability pairs may exhibit slight variations in shape. The tail heaviness of such distributions is determined by the parametric *kernel* distribution underlying the quantile parameterization. However, given the diverse range of QPDs proposed in the literature, a knowledgeable assessor should be able to select an appropriate distribution and validate the choice with the expert, taking into account the thickness of the distribution tails.

Most QPDs we reviewed are parameterized by a symmetric percentile triplet (SPT). These distributions rely on the symmetric property of underlying *kernel* distributions and can be generalized by swapping the distribution with another one that exhibits different tail shapes. Hadlock and Bickel (2019) utilized this method to generalize Johnson Quantile Parameterized distributions (J-QPDs). We show that the variants of Myerson distribution appearing in the literature (Myerson 2005, Wilson et al. 2023) represent similar generalization. This principle can be extended to include other kernels that result in varying thickness of the tails.

### 5.1. Quantile Function Perspective

The distributions discussed in this paper are defined using the quantile function, and therefore, they can be considered *quantile-based* quantile-parameterized distributions. Myerson, J-QPD, and several other quantile-parameterized distributions reparameterize conventional distributions, utilizing Gilchrist's transformations (Table 1).

Perepolkin et al. (2023) demonstrated that the distributions defined by quantile function can be used both as prior and as likelihood in Bayesian models. Priors defined by quantile function eliminate the need to compute prior density. The quantile function acts as a nonlinear transformation of a uniform degenerate random variate with the resulting Jacobian adjustment reciprocal to the density quantile function. Therefore, both the Jacobian and the density quantile function are

omitted from the Bayesian updating equation (Perepolkin et al. 2023). When using quantile-based QPDs as likelihood, special care needs to be taken with regard to the suitable prior for the QPP parameters. Perepolkin et al. (2024) used the Dirichet-based prior for the metalog likelihood model and described the *hybrid* elicitation process for encoding the expert judgments into the two-dimensional prior distribution implied by the model. Alternatively, Wilson et al. (2023) used multivariate normal distribution for the inter-quantile distances. This approach requires specification of the covariance structure, which might be difficult to elicit.

Not all QPDs are equally reliable in approximating the underlying distributions. Violating the QF transformation rules imposes additional constraints on the feasibility of parameters because certain combinations of parameters may result in locally decreasing quantile functions (Keelin 2016, Hadlock 2017).

## 5.2. Multivariate Extensions

Quantile-parameterized distributions can be readily extended to the multivariate setting by leveraging traditional multivariate distributions. The combination of quantile-based marginal distributions joined by the multivariate normal has been discussed previously in the literature (Hoff 2007, Drovandi and Pettitt 2011). Building on this approach, we proposed the use of Gumbel's bivariate logistic distribution (Gumbel 1961) to combine quantile-parameterized Logit-Myerson distributions (Wilson et al. 2023).

Copulas offer a natural extension of univariate QPDs into the multivariate domain. Bivariate copulas can be assembled into more complex structures using vine copulas (Kurowicka and Joe 2011, Wilson 2018, Czado 2019). Flexible QPDs serve as a viable alternative to empirical copulas, where the margins are represented by kernel density estimation (KDE) or other nonparametric approaches. Poorly fitted marginal distributions mean a *less-than-ideal* starting point for copula modeling because of potential deviations from uniformity of the copula margins.

Quantile-parameterized distributions defined by quantile function are particularly well-suited for constructing new distributions using bivariate quantiles (Vineshkumar and Nair 2019, Nair and Vineshkumar 2023). The ability to construct a conditional quantile

function as a linear combination of univariate quantile functions offers a convenient and interpretable approach to defining bivariate distributions, especially when the univariate quantile functions are parameterized by quantiles. These distributions are easy to sample from and construct. However, fitting these distributions to data can be challenging. As shown by Castillo et al. (1997) the fitting process requires all marginal and conditional quantile functions to be available in closed form, which is often unattainable.

## 5.3. Further Research

There appears to be a limited availability of unbounded quantile-parameterized distributions in the current literature. Among the distributions we examined, only the metalog distribution can extend across the entire real line. The G-QPD system provides clear distributional bounds defined explicitly by the expert during elicitation. In contrast, the (Generalized) Myerson distribution system relies on implicit bounds that need to be communicated to the expert. Most of the distributions we reviewed are characterized by a symmetrical percentile triplet (SPT) because they rely on the symmetrical property of their kernels. However, there may be situations where an arbitrary (nonsymmetrical) quantile parameterization could prove valuable (as shown by Perepolkin et al. 2024). The development of flexible quantile-parameterized distributions defined by an arbitrary set of quantile-probability pairs can enhance versatility of QPDs and facilitate their broader adoption.

In conclusion, quantile-parameterized distributions offer a valuable framework for capturing expert assessments and incorporating them into statistical models. They provide high flexibility and parameter interpretability, making them particularly beneficial for domain experts. The diverse range of quantile-parameterized distributions explored in the literature allows for customized modeling approaches that align with the expert's beliefs and uncertainties. By embracing these innovative distributions, researchers and practitioners can enhance the accuracy and reliability of their statistical models while leveraging expert knowledge effectively.

## Acknowledgments

The authors thank the editorial team and reviewers for their constructive feedback, which helped them improve this article.

## References

- Aas K, Czado C, Frigessi A, Bakken H (2009) Pair-copula constructions of multiple dependence. *Insurance Math. Econom.* 44(2):182–198.
- Akbarov A (2009) Probability elicitation: Predictive approach. PhD thesis, University of Salford.
- Arachchige CNPG, Prendergast LA, Staudte RG (2022) Robust analogs to the coefficient of variation. *J. Appl. Statist.* 49(2):268–290.
- Basikhasteh M, Lak F, Tahmasebi S (2021) Bayesian estimation of Morgenstern type bivariate Rayleigh distribution using some types of ranked set sampling. *Rev. Colomb. Estad.* 44(2):279–296.
- Berkson J (1951) Why I prefer logits to probits. *Biometrics* 7(4):327–339.
- Best N, Dallow N, Montague T (2020) Prior elicitation. Lesaffre E, Baio G, Boulanger B, eds. *Bayesian Methods in Pharmaceutical Research* (Chapman and Hall/CRC, Boca Raton, FL), 87–109.
- Bowley AL (1920) *Elements of Statistics* (Scribner's, New York).
- Burgman M, Layman H, French S (2021) Eliciting model structures for multivariate probabilistic risk analysis. *Front. Appl. Math. Statist.* 7(668037):1–10.
- Castillo E, Sarabia JM, Hadi AS (1997) Fitting continuous bivariate distributions to data. *J. Roy. Statist. Soc. D* 46(3):355–369.
- Chalabi Y, Scott DJ, Wuertz D (2012) Flexible distribution modeling with the generalized lambda distribution. Working Paper MPRA Paper No. 43333, ETH, Zurich, Switzerland.
- Cook J (2010) Determining distribution parameters from quantiles. Working paper, The University of Texas M. D. Anderson Cancer Center, Houston.
- Cullen AC, Frey HC, Frey CH (1999) *Probabilistic Techniques in Exposure Assessment: A Handbook for Dealing with Variability and Uncertainty in Models and Inputs* (Springer Science & Business Media, New York).
- Czado C (2019) *Analyzing Dependent Data with Vine Copulas* (Springer Berlin Heidelberg, New York).
- Drovandi CC, Pettitt AN (2011) Likelihood-free Bayesian estimation of multivariate quantile distributions. *Comput. Statist. Data Anal.* 55(9):2541–2556.
- Durrleman V, Nikeghbali A, Roncalli T (2000) A simple transformation of copulas. Preprint, submitted July 31, <http://dx.doi.org/10.2139/ssrn.1032543>.
- EFSA PoPH, Bragard C, Baptista P, Chatzivassiliou E, Di Serio F, Gonthier P, Jaques Miret JA, Justesen AF, et al. (2023) Risk assessment of *Citripestis sagittiferella* for the EU. *EFSA J.* 21(2):1–41.
- Elfadaly FG, Garthwaite PH (2017) Eliciting Dirichlet and Gaussian copula prior distributions for multinomial models. *Statist. Comput.* 27(2):449–467.
- Faber I, Jung J (2021) *Rmetalog: The Metalog Distribution* (CRAN).
- Field C, Genton MG (2006) The multivariate g-and-h distribution. *Technometrics* 48(1):104–111.
- Fiori AM, Zenga M (2009) Karl Pearson and the origin of kurtosis. *Internat. Statist. Rev.* 77(1):40–50.
- Freimer M, Kollia G, Mudholkar GS, Lin CT (1988) A study of the generalized Tukey lambda family. *Commun. Statist. Methods* 17(10):3547–3567.
- Genest C, Favre AC (2007) Everything you always wanted to know about copula modeling but were afraid to ask. *J. Hydrol. Engrg.* 12(4):347–368.
- Genest C, Quessy JF, Rémillard B (2006) Goodness-of-fit procedures for copula models based on the probability integral transformation. *Scandinavian J. Statist.* 33(2):337–366.
- Genest C, Rémillard B, Beaudoin D (2009) Goodness-of-fit tests for copulas: A review and a power study. *Insurance Math. Econom.* 44(2):199–213.
- Gilchrist W (2000) *Statistical Modelling with Quantile Functions* (Chapman & Hall/CRC, Boca Raton, FL).
- Gosling JP (2018) SHELFL: The Sheffield elicitation framework. Dias LC, Morton A, Quigley J, eds. *Elicitation* (Springer International Publishing, Cham, Switzerland), 61–93.
- Groeneveld RA (1998) A class of quantile measures for kurtosis. *Amer. Statist.* 52(4):325–329.
- Gumbel EJ (1960) Bivariate exponential distributions. *J. Amer. Statist. Assoc.* 55(292):698–707.
- Gumbel EJ (1961) Bivariate logistic distributions. *J. Amer. Statist. Assoc.* 56(294):335–349.
- Hadlock CC (2017) Quantile-parameterized methods for quantifying uncertainty in decision analysis. PhD thesis, University of Texas, Austin.
- Hadlock CC, Bickel JE (2017) Johnson quantile-parameterized distributions. *Decision Anal.* 14(1):35–64.
- Hadlock CC, Bickel JE (2019) The generalized Johnson quantile-parameterized distribution system. *Decision Anal.* 16(1):67–85.
- Hanea AM, Nane GF, Bedford T, French S, eds. (2021) *Expert Judgment in Risk and Decision Analysis*, International Series Operations Research Management Science, vol. 293 (Springer International Publishing, Cham, Switzerland).
- Hartmann M, Agiashvili G, Bürkner P, Klamí A (2020) Flexible prior elicitation via the prior predictive distribution. Preprint, submitted February 23, <https://arxiv.org/abs/2002.09868>.
- Haynes M, Mengersen K (2005) Bayesian estimation of g-and-k distributions using MCMC. *Comput. Statist.* 20(1):7–30.
- Haynes MA, MacGillivray HL, Mengersen KL (1997) Robustness of ranking and selection rules using generalised g-and-k distributions. *J. Statist. Plan. Inference* 65(1):45–66.
- Hemming V, Burgman MA, Hanea AM, McBride MF, Wintle BC (2018) A practical guide to structured expert elicitation using the IDEA protocol. *Methods Ecol. Evol.* 9(1):169–180.
- Herrerías-Velasco JM, Herrerías-Pleguezuelo R, van Dorp JR (2009) The generalized two-sided power distribution. *J. Appl. Statist.* 36(5):573–587.
- Hoff PD (2007) Extending the rank likelihood for semiparametric copula estimation. *Ann. Appl. Statist.* 1(1):265–283.
- Holzhauser B, Hampson LV, Gosling JP, Bornkamp B, Kahn J, Lange MR, Luo WL, et al. (2022) Eliciting judgements about dependent quantities of interest: The SHEffield ELicitation Framework extension and copula methods illustrated using an asthma case study. *Pharm. Statist.* 21(5):1005–1021.
- Ingram B (2020) *Rjqpd: The Johnson Quantile-Parameterised Distribution* (CRAN).
- Jeong-Soo P (2005) Wakeby distribution and the maximum likelihood estimation algorithm in which probability density function is not explicitly expressed. *Commun. Statist. Appl. Methods* 12(2): 443–451.
- Johnson D (1997) The triangular distribution as a proxy for the beta distribution in risk analysis. *J. Roy. Statist. Soc. D* 46(3):387–398.
- Johnson NL, Kotz S, Balakrishnan N (1994) *Continuous Univariate Distributions*. Wiley Series in Probability and Mathematical Statistics, 2nd ed. (Wiley, New York).
- Jones MC, Rosco JF, Pewsey A (2011) Skewness-invariant measures of kurtosis. *Amer. Statist.* 65(2):89–95.
- Kadane JB (1980) Predictive and structural methods for eliciting prior distributions. Zellner A, ed. *Bayesian Analysis in Econometrics and Statistics: Essays in Honor of Harold Jeffreys* (North Holland Publishing Company, Amsterdam), 89–93.

- Kadane J, Wolfson LJ (1998) Experiences in elicitation. *J. Roy. Statist. Soc. Ser. D Statist.* 47(1):3–19.
- Keelin TW (2016) The metalog distributions. *Decision Anal.* 13(4): 243–277.
- Keelin TW (2017) The metalog distributions—feasibility. Accessed July 3, 2024, <http://www.metalogdistributions.com/equations/feasibility.html>.
- Keelin TW, Howard RA (2021) The metalog distributions: Virtually unlimited shape flexibility, combining expert opinion in closed form, and Bayesian updating in closed form. Preprint, submitted October 12, <https://osf.io/xdg5e/>.
- Keelin TW, Powley BW (2011) Quantile-parameterized distributions. *Decision Anal.* 8(3):206–219.
- Khan I (2020) Jqpd: The python JQPD. Accessed July 3, 2024, <https://pypi.org/project/jqpd/>.
- Kim TH, White H (2004) On more robust estimation of skewness and kurtosis. *Finance Res. Lett.* 1(1):56–73.
- Kotz S, Van Dorp JR (2004) *Beyond Beta: Other Continuous Families of Distributions with Bounded Support and Applications* (World Scientific, Singapore/Hackensack, NJ).
- Kotz S, Van Dorp JR (2005) A link between two-sided power and asymmetric Laplace distributions: With applications to mean and variance approximations. *Statist. Probab. Lett.* 71(4):383–394.
- Kurowicka D, Cooke R (2006) *Uncertainty Analysis with High Dimensional Dependence Modelling*. Wiley Series in Probability and Statistics (Wiley, Chichester, UK/Hoboken, NJ).
- Kurowicka D, Joe H, eds. (2011) *Dependence Modeling: Vine Copula Handbook* (World Scientific, Singapore).
- Lampasi D (2008) An alternative approach to measurement based on quantile functions. *Measurement* 41(9):994–1013.
- Mac Gillivray H (1992) Shape properties of the g-and-h and Johnson families. *Commun. Statist. Theory Methods* 21(5):1233–1250.
- Mikkola P, Martin OA, Chandramouli S, Hartmann M, Pla OA, Thomas O, Pesonen H, et al. (2023) Prior knowledge elicitation: The past, present, and future. *Bayesian Anal.* 1(1):1–33.
- Moors JJA (1988) A quantile alternative to kurtosis. *J. Roy. Statist. Soc. Ser. D Statist.* 37(1):25–32.
- Morgan MG (2014) Use (and abuse) of expert elicitation in support of decision making for public policy. *Proc. Natl. Acad. Sci. USA* 111(20):7176–7184.
- Myerson RB (2005) *Probability Models for Economic Decisions*. Duxbury Applied Series (Thomson/Brooke/Cole, Belmont, CA).
- Nair NU, Vineshkumar B (2023) Properties of bivariate distributions represented through quantile functions. *Amer. J. Math. Management Sci.* 42(1):1–12.
- O'Hagan A (2019) Expert knowledge elicitation: Subjective but scientific. *Amer. Statist.* 73(sup1):69–81.
- O'Hagan A, Buck CE, Daneshkhah A, Eiser JR, Garthwaite PH, Jenkinson DJ, Oakley JE, Rakow T (2006) *Uncertain Judgements: Eliciting Experts' Probabilities: O'Hagan/Uncertain Judgements: Eliciting Experts' Probabilities* (John Wiley & Sons, Chichester, UK).
- Palisade Corp (2009) Guide to using@ RISK.: Risk analysis and simulation add-in for Microsoft Excel. Technical report, Palisade Corporation, Newfield, NY.
- Parzen E (1979) Nonparametric statistical data modeling. *J. Amer. Statist. Assoc.* 74(365):105–121.
- Perepolkin D (2019) *Qpd: Tools for Quantile-Parameterized Distributions*. Accessed July 3, 2024, <https://github.com/dmi3kno/qpd>.
- Perepolkin D, Goodrich B, Sahlin U (2023) The tenets of quantile-based inference in Bayesian models. *Comput. Statist. Data Anal.* 187(107795):1–15.
- Perepolkin D, Goodrich B, Sahlin U (2024) Hybrid elicitation and quantile-parameterized likelihood. *Statist. Comput.* 34(11):1–15.
- Powley BW (2013) Quantile function methods for decision analysis. PhD thesis, Stanford University, Palo Alto, CA.
- Prangle D (2017) gk: An R package for the g-and-k and generalised g-and-h distributions. Preprint, submitted June 21, <https://arxiv.org/abs/1706.06889>.
- Rahman A, Zaman MA, Haddad K, El Adlouni S, Zhang C (2015) Applicability of Wakeby distribution in flood frequency analysis: A case study for eastern Australia. *Hydrol. Processes* 29(4): 602–614.
- Ramberg JS, Schmeiser BW (1974) An approximate method for generating asymmetric random variables. *Commun. ACM* 17(2):78–82.
- Rayner GD, MacGillivray HL (2002) Numerical maximum likelihood estimation for the g-and-k and generalized g-and-h distributions. *Statist. Comput.* 12(1):57–75.
- Sajeevkumar N, Irshad M (2014) Estimation of a parameter of Morgenstern type bivariate logistic distribution with equal coefficients of variation by concomitants of order statistics. *Calcutta Statist. Assoc. Bull.* 66(3–4):213–228.
- Sharma R, Das S (2018) Regularization and variable selection with copula prior. Preprint, submitted September 16, <https://arxiv.org/abs/1709.05514>.
- Shih N (2015) The model identification of beta distribution based on quantiles. *J. Statist. Comput. Simul.* 85(10):2022–2032.
- Smith MS (2013) Bayesian approaches to copula modelling. Preprint, submitted December 18, <https://arxiv.org/abs/1112.4204>.
- Smith C, Jefferies T, Faber I (2021) Pymetalog: The python metalog distribution. <https://github.com/tjefferies/pymetalog>.
- Spetzler CS, Staël Von Holstein CAS (1975) Probability encoding in decision analysis. *Management Sci.* 22(3):340–358.
- Tarsitano A (2005) Fitting Wakeby model using maximum likelihood. *Statistica e Ambiente* 1(1):253–256.
- Tukey JW (1965) Which part of the sample contains the information? *Proc. Natl. Acad. Sci. USA* 53(1):127–134.
- van Dorp JR, Mazzuchi TA (2000) Solving for the parameters of a beta a distribution under two quantile constraints. *J. Statist. Comput. Simul.* 67(2):189–201.
- van Dorp JR, Mazzuchi TA (2021) Three-point lifetime distribution elicitation for maintenance optimization in a Bayesian context. Hanea A, Nane GF, Bedford T, French S, eds. *Expert Judgement in Risk and Decision Analysis*, International Series in Operations Research & Management Science (Springer International Publishing, Cham, Switzerland), 147–177.
- van Dorp JR, Shittu E (2023) The generalized two-sided beta distribution with applications in project risk analysis. *J. Statist. Theory Pract.* 17(38):1–27.
- Vineshkumar B, Nair NU (2019) Bivariate quantile functions and their applications to reliability modelling. *Statistics* 79(1):3–21.
- Wang W, Wells MT (2000) Model selection and semiparametric inference for bivariate failure-time data. *J. Amer. Statist. Assoc.* 95(449):62–72.
- Welsh MB, Begg SH (2018) More-or-less elicitation (MOLE): Reducing bias in range estimation and forecasting. *EURO J. Decision Process* 6(1):171–212.

- Wilson KJ (2018) Specification of informative prior distributions for multinomial models using vine copulas. *Bayesian Anal.* 13(3): 749–766.
- Wilson KJ, Elfadaly FG, Garthwaite PH, Oakley JE (2021) Recent advances in the elicitation of uncertainty distributions from experts for multinomial probabilities. Hanea AM, Nane GF, Bedford T, French S, eds. *Expert Judgement in Risk and Decision Analysis*, International Series in Operations Research & Management Science (Springer International Publishing, Cham, Switzerland), 19–51.
- Wilson KJ, Farrow M, French S, Hartley D (2023) Reconciliation of expert priors for quantities and events and application within the probabilistic Delphi method. Preprint, submitted November 24, <https://arxiv.org/abs/2311.14487>.
- Winkler RL (1980) Prior information, predictive distributions, and Bayesian model-building. Zellner A, ed. *Bayesian Analysis in Econometrics and Statistics: Essays in Honor of Harold Jeffreys (Studies in Bayesian Econometrics)* (North-Holland Publishing Company, Amsterdam), 95–109.

---

**Dmytro Perepolkin** is a recent PhD graduate in Environmental Science from Lund University, driven by a passion

for teaching data literacy and introductory Bayesian analysis at the university level. His research explores applications of the quantile function approach to traditional statistical challenges, the integration of expert judgment in Bayesian inference, and the philosophical underpinnings of structured decision-making to promote rationality.

**Erik Lindström** is a professor of mathematical statistics at Lund University, specializing in statistical methodology, financial mathematics, and energy finance. His research spans time series analysis, energy market stability, and algorithm development for parameter estimation in continuous-time processes. He earned his PhD in mathematical statistics from Lund University.

**Ullrika Sahlin** is an associate professor in Environmental Science at Lund University. Her research focuses on characterizing uncertainty in scientific assessment using subjective probability and Bayesian inference, with applications on assessment of human health risk, biodiversity, and other ecological endpoints. She is interested in developing methodologies and guidance in the overlap between complex statistical modeling, expert knowledge elicitation, and uncertainty analysis.